

A MARTINGALE INEQUALITY FOR THE SQUARE  
AND MAXIMAL FUNCTIONS

BY

LOUIS H.Y. CHEN

TECHNICAL REPORT NO. 31

MARCH 15, 1979

PREPARED UNDER GRANT

DAAG29-77-G-0031

FOR THE U.S. ARMY RESEARCH OFFICE

Reproduction in Whole or in Part is Permitted  
for any purpose of the United States Government  
Approved for public release; distribution unlimited.

DEPARTMENT OF STATISTICS  
STANFORD UNIVERSITY  
STANFORD, CALIFORNIA



A MARTINGALE INEQUALITY FOR THE SQUARE  
AND MAXIMAL FUNCTIONS

By

LOUIS H. Y. CHEN

TECHNICAL REPORT NO. 31  
MARCH 15, 1979

Prepared under Grant DAAG29-77-G-0031  
For the U.S. Army Research Office  
Herbert Solomon, Project Director

Approved for public release; distribution unlimited.

DEPARTMENT OF STATISTICS  
STANFORD UNIVERSITY  
STANFORD, CALIFORNIA

Partially supported under Office of Naval Research Contract N00014-76-C-0475  
(NR-042-267) and issued as Technical Report No. 270.

The findings in this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

# A MARTINGALE INEQUALITY FOR THE SQUARE

## AND MAXIMAL FUNCTIONS

By

Louis H. Y. Chen

University of Singapore and Stanford University

### 1. Introduction and Notation.

An inequality involving a class of functions for weak martingales and nonnegative weak submartingales is proved. Three special cases are deduced, one of which generalizes and refines a result of Austin (1966). As an application of the inequality, the special cases are used to give easy proofs of Burkholder's (1966)  $L \log L$  and  $L_p$  (for  $1 < p \leq 2$ ) inequalities for the square function of a martingale or a nonnegative submartingale. Although the inequality and the special cases are proved for weak martingales and nonnegative weak submartingales, they are also new for martingales and nonnegative submartingales.

Weak martingales and weak submartingales were first defined in Nelson (1970). Examples of these can be found in Nelson (1970) and Berman (1976). For ease of reference, we give the definition here. A sequence  $f = (f_1, f_2, \dots)$  defined on a probability space is a weak martingale (or weak submartingale) if  $f_n$  is integrable and  $E(f_{n+1} | f_n) = (\text{or } \geq) f_n$  a.s. for  $n \geq 1$ . Of course, a martingale (or submartingale) is a weak martingale (or weak submartingale).

Throughout this paper, unless otherwise stated,  $f = (f_1, f_2, \dots)$  will denote a weak martingale or a nonnegative weak submartingale. As usual  $f_0 \equiv 0$ . The difference sequence of  $f$  will be denoted by

$d = (d_1, d_2, \dots)$ . Also  $S_n(f) = \left( \sum_{i=1}^n d_i^2 \right)^{1/2}$ ,  $S(f) = \sup_{1 \leq n < \infty} S_n(f)$ ,  
 $f_n^* = \sup_{1 \leq i \leq n} |f_i|$ ,  $f^* = \sup_{1 \leq n < \infty} f_n^*$  and  $\|f\|_p = \sup_{1 \leq n < \infty} \|f_n\|_p$  for  
 $1 \leq p < \infty$ . All functions are real-valued, Borel measurable and  
defined on the real line.

## 2. The Main Results.

We first derive an identity.

Lemma 2.1. Let  $\varphi$  be a differentiable function whose derivative  $\varphi'$  is an indefinite integral of  $\varphi''$  such that  $\varphi(0) = \varphi'(0) = 0$ ,  $\varphi''$  is a nonnegative and even function, and such that for  $n \geq 1$ ,  $f_n \varphi'(f_n)$  is integrable. Define  $K_i(x) = (d_i - x)^+$  if  $x \geq 0$  and  $= (d_i - x)^-$  if  $x < 0$ ,  $i \geq 1$ . Then for  $n \geq 1$ ,  $\varphi(f_n)$  is integrable, and

$$(2.1) \quad E\varphi(f_n) \geq \sum_{i=1}^n E \int_{-\infty}^{\infty} \varphi''(f_{i-1} + x) K_i(x) dx$$

where equality holds in the weak martingale case. Furthermore for  $i \geq 1$ .

$$(2.2) \quad \int_{-\infty}^{\infty} K_i(x) dx = \frac{1}{2} d_i^2.$$

Proof. Since the proof of (2.2) is easy, we omit it here. Since  $\varphi(0) = \varphi'(0) = 0$ , we have  $\varphi'(x) = \int_0^x \varphi''(t) dt$  and  $\varphi(x) = \int_0^x \varphi'(t) dt$ . It follows that  $\varphi'$  is an odd function and  $\varphi$  an even function. Therefore

$$(2.3) \quad 0 \leq \varphi(x) = \int_0^{|x|} \varphi'(t) dt = |x| \varphi'(|x|) - \int_0^{|x|} t \varphi''(t) dt \\ \leq x \varphi'(x) .$$

The integrability of  $\varphi(f_n)$  then follows from that of  $f_n \varphi'(f_n)$ . We also need the integrability of  $d_i \varphi'(f_{i-1})$  for  $i \geq 1$ . Since  $\varphi'' \geq 0$ ,  $\varphi'$  is nondecreasing. Therefore

$$|f_i| \varphi'(|f_{i-1}|) I(|f_{i-1}| \leq |f_i|) \leq |f_i| \varphi'(|f_i|)$$

and

$$|f_i| \varphi'(|f_{i-1}|) I(|f_{i-1}| > |f_i|) \leq |f_{i-1}| \varphi'(|f_{i-1}|) .$$

Hence

$$|d_i \varphi'(f_{i-1})| = |d_i| \varphi'(|f_{i-1}|) \\ \leq |f_i| \varphi'(|f_{i-1}|) + |f_{i-1}| \varphi'(|f_{i-1}|) \\ \leq |f_i| \varphi'(|f_i|) + 2|f_{i-1}| \varphi'(|f_{i-1}|) .$$

This implies the integrability of  $d_i \varphi'(f_{i-1})$ . We now derive (2.1).

In the weak martingale case, the left hand side of (2.1) is equal to

$$(2.4) \quad \sum_{i=1}^n E[\varphi(f_i) - \varphi(f_{i-1})] \\ = \sum_{i=1}^n E[\varphi(f_i) - \varphi(f_{i-1}) - d_i \varphi'(f_{i-1})] \\ = \sum_{i=1}^n E \left\{ \int_0^{d_i} \int_0^y \varphi''(f_{i-1}+x) dx dy \right\} I(d_i \geq 0) \\ + \sum_{i=1}^n E \left\{ \int_{d_i}^0 \int_y^0 \varphi''(f_{i-1}+x) dx dy \right\} I(d_i < 0) .$$

In the nonnegative weak submartingale case,  $\varphi'(f_{i-1})$  is nonnegative and (2.4) holds with the first equality replaced by " $\geq$ ". Now  $\varphi'' \geq 0$ . So we may reverse the order of the doubly integration in (2.4). By this, the extreme right hand side of (2.4) yields

$$\sum_{i=1}^n E \int_{-\infty}^{\infty} \varphi''(f_{i-1}+x) K_i(x) dx$$

and the lemma is proved.

In the case where  $f$  is a martingale or a nonnegative submartingale, let  $\tau$  be a stopping time. By replacing  $f$  in (2.1) by the stopped martingale or nonnegative submartingale  $f^\tau$ , we obtain

$$(2.5) \quad E\varphi(f_n^\tau) \geq \sum_{i=1}^n E I(\tau \geq i) \int_{-\infty}^{\infty} \varphi''(f_{i-1}+x) K_i(x) dx$$

where again equality holds in the martingale case. If the differences of  $f$  are mutually independent with zero means,  $\varphi(x) = x^2$  and  $\tau = \inf\{n: |f_n| > a\}$  where  $a > 0$ , then (2.5) immediately yields Kolmogorov's two inequalities in the proof of the three series theorem.

Theorem 2.1. Let  $\psi'$  be a nonnegative and even function which is nonincreasing on  $[0, \infty)$ , and let  $\psi(x) = \int_0^x \psi'(t) dt$ . Then

$$(2.6) \quad ES^2(f) \psi'(f^*) \leq 2 \sup_n E |f_n| \psi(|f_n|).$$

Proof. There is nothing to prove if the right hand side of (2.6) is infinite. So we assume it to be finite. Let  $K_i(x)$  be as in Lemma 2.1. It is not difficult to see that, for  $i \geq 1$ ,  $f_{i-1}+x$  lies

between  $f_{i-1}$  and  $f_i$  on  $\{x: K_i(x) > 0\}$ . Now let  $\varphi'' = \psi'$ ,  
 $\varphi'(x) = \int_0^x \varphi''(t)dt = \psi(x)$  and  $\varphi(x) = \int_0^x \varphi'(t)dt$ . Then the integrability condition in Lemma 2.1 is satisfied and the lemma immediately yields

$$E S_n^2(f) \psi'(f_n^*) \leq 2E\varphi(f_n) \leq 2E|f_n| \psi(|f_n|) \leq 2 \sup_n E|f_n| \psi(|f_n|)$$

where the second inequality follows from (2.3). By letting  $n \rightarrow \infty$  and applying Fatou's lemma, the theorem is proved.

We now deduce from (2.6) three special cases.

Corollary 2.1. We have

$$(2.7) \quad E \frac{S^2(f)}{1+f^{*2}} \leq \pi \|f\|_1 ;$$

$$(2.8) \quad E \frac{S^2(f)}{1+f^*} \leq 2 \sup_n E|f_n| \log(1+|f_n|) ;$$

$$(2.9) \quad E \frac{S^2(f)}{f^{*2-p}} \leq \frac{2}{p-1} \|f\|_p^p, \quad 1 < p \leq 2 .$$

Proof. For (2.7), let  $\psi'(x) = (1+x^2)^{-1}$ ; and for (2.8), let  $\psi'(x) = (1+|x|)^{-1}$ . For (2.9), we first let  $\psi'(x) = (a+|x|)^{p-2}$  where  $a > 0$  and then let  $a \downarrow 0$ .

The inequality (2.7) generalizes and refines a result of Austin (1966) who proved that the square function of an  $L_1$ -bounded martingale is square integrable on any set where the maximal function is bounded above.



### 3. Applications.

In this section, we use Corollary 2.1 to give easy proofs of Burkholder's  $L \log L$  and  $L_p$  (for  $1 < p \leq 2$ ) inequalities for the square function of a martingale or a nonnegative submartingale. These inequalities were first proved by Burkholder (1966). Since then different proofs have been given. (See, for example, Gordon (1972), Burkholder (1973), Chao (1973) and Garsia (1973).)

Theorem 3.1. Let  $f = (f_1, f_2, \dots)$  be a martingale or a nonnegative submartingale. Then

$$(3.1) \quad ES(f) \leq 2 \left( \frac{e}{e-1} \right)^{1/2} [1 + \sup_n E|f_n| \log^+ |f_n|] .$$

Proof. We shall use the following inequality which dates back to Young (1913). It can also be found in Doob (1953).

$$(3.2) \quad a \log^+ b \leq a \log^+ a + be^{-1} \quad \text{for } a \geq 0 \text{ and } b \geq 0.$$

Replacing  $f_i$  by  $\lambda^{-1} f_i$  in (2.8) of Corollary 2.1 where  $\lambda = Ef^*$ , we obtain

$$(3.3) \quad E \frac{S^2(f)}{\lambda + f^*} \leq 2 \sup_n E|f_n| \log(1 + \lambda^{-1} |f_n|)$$

which by (3.2)

$$\leq 2 \sup_n [E|f_n| \log^+ |f_n| + (\lambda e)^{-1} (\lambda + E|f_n|)]$$

$$\leq 2[1 + \sup_n E|f_n| \log^+ |f_n|] .$$

Now applying the Cauchy-Schwarz inequality to

$$ES(f) = E\left(\frac{S^2(f)}{\lambda + f^*}\right)^{1/2} (\lambda + f^*)^{1/2}$$

and using (3.3) and the following inequality of Doob (1953) for submartingales,

$$\|f^*\|_1 \leq \left(\frac{e}{e-1}\right) [1 + \sup_n E|f_n| \log^+ |f_n|] ,$$

we obtain (3.1). This proves the theorem.

Theorem 3.2. Let  $f = (f_1, f_2, \dots)$  be a martingale or a nonnegative submartingale. Then for  $1 < p \leq 2$ ,

$$(3.4) \quad \|S(f)\|_p \leq 2^{1/2} p^{1/2} q \|f\|_p$$

where  $p^{-1} + q^{-1} = 1$ .

Proof. Applying Hölder's inequality to

$$\|S(f)\|_p^p = E\left(\frac{S^2(f)}{f^{*2-p}}\right)^{\frac{1}{2}p} (f^{*p})^{1 - \frac{1}{2}p} ,$$

we obtain

$$\|S(f)\|_p^p \leq \left(E \frac{S^2(f)}{f^{*2-p}}\right)^{\frac{1}{2}p} (\|f^*\|_p^p)^{1 - \frac{1}{2}p}$$

which by (2.9) of Corollary 2.1

$$\leq \left(\frac{2}{p-1} \|f\|_p^p\right)^{\frac{1}{2}p} (\|f^*\|_p^p)^{1 - \frac{1}{2}p} .$$

This together with the following inequality of Doob (1953) for sub-martingales,

$$\|f^*\|_p \leq q \|f\|_p \quad \text{for } 1 < p < \infty, \quad p^{-1} + q^{-1} = 1,$$

imply

$$\|S(f)\|_p \leq \left(\frac{2}{p-1}\right)^{1/2} q^{1 - \frac{1}{2}p} \|f\|_p \leq 2^{1/2} p^{1/2} q \|f\|_p.$$

This proves the theorem.

The absolute constants in (3.1) and (3.4) seem to be the lowest ever obtained. In (3.4), the order of magnitude of the constant as  $p \rightarrow 1$  is the same as that obtained by Burkholder (1973).

## References

1. Austin, D. G. (1966). A sample function property of martingales. Ann. Math. Statist., 37, 1396-1397.
2. Berman, Jeremy (1976). An example of a weak martingale. Ann. Probability 4, 107-108.
3. Burkholder, D. L. (1966). Martingale transforms. Ann. Math. Statist., 37, 1494-1504.
4. Burkholder, D. L. (1973). Distribution function inequalities for martingales. Ann. Probability 1, 19-42.
5. Chao, Jia-Arng (1973). A note on martingale square functions. Ann. Probability 1, 1059-1060.
6. Doob, J. L. (1953). Stochastic Processes. Wiley, New York.
7. Garsia, Adriano M. (1973). Martingale Inequalities: Seminar Notes on Recent Progresses. Benjamin, Reading, Massachusetts.
8. Gordon, Louis (1972). An equivalent to the martingale square function inequality. Ann. Math. Statist., 43, 1927-1934.
9. Nelson, Paul I. (1970). A class of orthogonal series related to martingales. Ann. Math. Statist., 41, 1684-1694.
10. Young, W. H. (1913). On a certain series of Fouries. Proc. London Math. Soc., 11, 357-366.

Department of Mathematics  
University of Singapore  
Bukit Timah Road  
Singapore 10

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 31	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A MARTINGALE INEQUALITY FOR THE SQUARE AND MAXIMAL FUNCTIONS		5. TYPE OF REPORT & PERIOD COVERED TECHNICAL REPORT
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) LOUIS H. Y. CHEN		8. CONTRACT OR GRANT NUMBER(s) DAAG29-77-G-0031
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics Stanford University Stanford, CA 94305		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS P-14435-M
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office Post Office Box 12211 Research Triangle Park, NC 27709		12. REPORT DATE MARCH 15, 1979
		13. NUMBER OF PAGES 9
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES The findings in this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents. This report partially supported under Office of Naval Research Contract N00014-76-C-0475 (NR-042-267) and issued as Technical Report No. 270.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Martingale inequality; Maximal function; Square function; Weak martingale.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  PLEASE SEE REVERSE SIDE		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

S/N 0102-LF-014-6691

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

## A Martingale Inequality for the Square and Maximal Functions

An inequality for weak martingales and nonnegative weak submartingales is proved. Three special cases are deduced, one of which generalizes and refines a result of Austin (1966). As an application of the inequality, the special cases are used to give easy proofs of Burkholder's (1966)  $L \log L$  and  $L_p$  (for  $1 < p \leq 2$ ) inequalities for the square function of a martingale or a nonnegative submartingale.

270/31 - Author: Louis H.Y. Chen